

BEST TECHNOLOGY ■ BROAD-SPECTRUM WEED CONTROL ■ STRONG STEWARDSHIP

You have the Power to Control what You Grow, Including Red Rice

 **BASF**
The Chemical Company



CLEARFIELD[®]
production system | rice

Residual Weed Control with Multiple Modes of Action


CLEARFIELD® technology delivers more than just weed control. It effectively controls tough broadleaf and grass weeds like red rice which can translate into increased yields and cleaner grain at harvest; all with simplicity, convenience and improved return on investment.

Residual CLEARFIELD® Rice Herbicide Program

The first application, **Clearpath**® herbicide, should be applied between emergence and 2nd leaf. The second application, **Newpath**® herbicide, should be applied approximately 2 weeks later between the 3 and 5-leaf stages (prior to tillering).

Beyond® herbicide may be used as a single rescue treatment for weed breakthroughs any time following the **Newpath** application and prior to Panicle Initiation (or Green Ring).

Facet® herbicide may be applied as a soil treatment, after emergence, or between the 2-leaf stage and permanent flood. No more than 0.67 lb/A of **Facet** should be applied per season.



Note: **Beyond** may be applied up to 14 days following Green Ring or Panicle Initiation in **CLEARFIELD** rice varieties.



Untreated



Clearpath (0.5 lb/A) followed by **Newpath** (4 oz/A)

Knockdown and Control Tough, Late-season Weeds


Now, the **CLEARFIELD**® Production System for rice is more flexible with a broadened **CLEARFIELD** herbicide application window to target tough-to-control, late season weeds; all in an effort to help you get the most out of every rice acre on your farm.

Knockdown CLEARFIELD® Rice Herbicide Program

Clearpath® herbicide should be applied to rice from emergence through 5-leaf.

Beyond® herbicide should be applied approximately 2 weeks later, between the 4 to 5-leaf stages.

Beyond can also be applied *again* as a rescue treatment prior to Panicle Initiation (or Green Ring)



Note: **Beyond** may be applied up to 14 days following Green Ring or Panicle Initiation in **CLEARFIELD** rice varieties.

Stewardship Recommendations:

- Start clean - utilize a burndown and residual herbicide at planting
- Do not plant **CLEARFIELD** rice in consecutive years
- Rotate herbicides with alternative modes of action
- Read and follow **CLEARFIELD** Rice Stewardship Guidelines and Agreement

Get the Most Out of Every CLEARFIELD® Rice Acre

Brand	Active Ingredient	AI Load	Use Rate	Timing	Adjuvant	Target Weed(s)
Prowl® H₂O Herbicide An important tool for resistance management, Prowl H₂O also allows you to delay flooding with its residual control of tough grasses like barnyardgrass, panicum, sprangletop and broadleaf weeds including pigweed spp.	Pendimethalin	3.8 lb/gal	2 pt/A	Delayed preemerge to early postemerge	-	junglerice, sprangletop and barnyardgrass
Clearpath® Herbicide The most powerful and convenient combination of the same active ingredients found in Newpath® herbicide and Facet® herbicide designed to deliver unsurpassed, broad-spectrum weed control and resistance management delivering two modes of action.*	Imazethapyr + Quinclorac	75% Combined	0.5 to 0.72 lb/A	Early postemerge prior to 5-leaf rice stage (dry-seeded)	COC (2 pt/A)	red rice, barnyardgrass, broadleaf signalgrass and hemp sesbania
Newpath Herbicide The industry standard providing residual and broad-spectrum weed control in the fight against red rice, barnyardgrass and other tough weeds in fields where CLEARFIELD® rice is grown.*	Imazethapyr	2 lb/gal	4 to 6 oz/A	Preplant, preemerge or early postemerge prior to 5-leaf rice stage	COC (1% v/v)	red rice and barnyardgrass
Facet 75DF Herbicide Provides reliable and consistent broad-spectrum control of broadleaf weeds and grasses, such as hemp sesbania, joint-vetches and propanil-resistant barnyardgrass.	Quinclorac	75%	0.4 to 0.67 lb/A	Early postemerge or spike at 2-leaf rice stage	COC (2 pt/A)	barnyardgrass, broadleaf signalgrass and hemp sesbania
Beyond® Herbicide A unique, flexible and broad-spectrum tool powered to knockdown advanced, late-season weeds including barnyardgrass, fall panicum and red rice.	Imazamox	1 lb/gal	5 to 6 oz/A	Early postemerge 14-21 days after Clearpath or Newpath application but before panicle initiation	COC (1% v/v)	red rice, barnyardgrass, broadleaf signalgrass

*For use only on CLEARFIELD rice varieties and hybrids (not less than 75% hybrid seed)

Always read and follow label directions.

CLEARFIELD, Beyond, Clearpath, Facet, Newpath and Prowl H₂O are registered trademarks of BASF. ©2010 BASF Corporation. All rights reserved. APN-2010-19-175-CRB-004

For more information on BASF Crop Protection products, visit www.agproducts.basf.us



CLEARFIELD®
production system | rice